

Dictyostelium discoideum and autophagy – a perfect pair

SARAH FISCHER and LUDWIG EICHINGER*

Centre for Biochemistry, Institute of Biochemistry I, Medical Faculty, University of Cologne, Cologne, Germany

ABSTRACT Autophagy is subdivided into chaperone-mediated autophagy, microautophagy and macroautophagy and is a highly conserved intracellular degradative pathway. It is crucial for cellular homeostasis and also serves as a response to different stresses. Here we focus on macroautophagy, which targets damaged organelles and large protein assemblies, as well as pathogenic intracellular microbes for destruction. During this process, cytosolic material becomes enclosed in newly generated double-membrane vesicles, the so-called autophagosomes. Upon maturation, the autophagosome fuses with the lysosome for degradation of the cargo. The basic molecular machinery that controls macroautophagy works in a sequential order and consists of the ATG1 complex, the PtdIns3K complex, the membrane delivery system, two ubiquitin-like conjugation systems, and autophagy adaptors and receptors. Since the different stages of macroautophagy from initiation to final degradation of cargo are tightly regulated and highly conserved across eukaryotes, simple model organisms in combination with a wide range of techniques contributed significantly to advance our understanding of this complex dynamic process. Here, we present the social amoeba *Dictyostelium discoideum* as an advantageous and relevant experimental model system for the analysis of macroautophagy.

KEY WORDS: *autophagy, ubiquitin-proteasome system (UPS), LC3-associated phagocytosis, proteaphagy*

Introduction

Macroautophagy, hereafter denoted as autophagy for simplicity, is the major lysosomal route for the clearance and turnover of damaged organelles and long-lived proteins (Stanley *et al.*, 2014). This cellular “self-eating” phenomenon was discovered in the sixties of the last century and the term “autophagy” was already coined in 1963 (De Duve and Wattiaux, 1966). Research in this process started slow, but is nowadays booming and increasingly fascinating. This was underlined by the award of the 2016 Nobel Prize in Physiology or Medicine to Yoshinori Ohsumi for his fundamental discoveries on the autophagic machinery in the budding yeast *Saccharomyces cerevisiae* (Levine and Klionsky, 2017, Tsukada and Ohsumi, 1993). Autophagy occurs in all eukaryotes at a basal level and is induced in response to cellular stresses such as starvation, the presence of protein aggregates or invading pathogens (Eskelinen and Saftig, 2009, Mizushima *et al.*, 2008). Most likely, autophagy evolved in unicellular organisms as a survival mechanism during starvation through the recycling of cellular building blocks (Wirawan *et al.*, 2012). The different molecular complexes act in a sequential way to deliver cytoplasmic cargo to the lysosome and crucial components were initially discovered and characterised in *S. cerevisiae* (Tsukada and Ohsumi, 1993). The hallmark of

autophagy is the *de novo* formation of a double-layered vesicle, the so-called autophagosome, which can engulf parts of the cytoplasm, entire organelles or even pathogenic bacteria (Lamb *et al.*, 2013). The maturation of autophagosomes into autolysosomes is accomplished by fusion of the outer autophagosomal membrane with the lysosomal membrane. Then, the inner autophagosomal membrane and the cargo are degraded by lysosomal hydrolases (Stanley *et al.*, 2014). The proteins involved in autophagosome formation were named ATG, for AuTophagy-related proteins, and are evolutionarily highly conserved across the eukaryotic lineage (Feng *et al.*, 2014). Autophagic dysfunction can result in a wide range of diseases, including neurodegeneration, cancer, muscular dystrophy, and lipid-storage disorders (Mizushima *et al.*, 2008, Schneider and Cuervo, 2014).

Autophagy has long been considered as a non-selective bulk degradation process of cytoplasmic components in response to nutrient starvation (Boya *et al.*, 2013). However, it is now clear that autophagy generally operates in a selective manner through recognition of the substrates by cargo specific autophagy recep-


Abbreviations used in this paper: LAP, LC3-associated phagocytosis; UPS, ubiquitin-proteasome system.

*Address correspondence to: Ludwig Eichinger. Centre for Biochemistry, Institute of Biochemistry I, Medical Faculty, University of Cologne, Cologne, Germany. Tel.: +49 221 478 6928; Fax: +49 221 478 97524. E-mail: ludwig.eichinger@uni-koeln.de -  <https://orcid.org/0000-0003-1594-6117>

Submitted: 29 May, 2019; Accepted: 9 July, 2019.

Fig. 1. Schematic depiction of autophagosome formation in *D. discoideum*.

During autophagy, cytosolic material becomes engulfed by the cup-shaped isolation membrane, which elongates and closes to form a double-membrane autophagosome. After fusion with a lysosome, the cargo is degraded and recycled. Different components are mandatory for this process: the ATG1 complex, the PtdIns3K complex, the membrane delivery system and two ubiquitin-like conjugation systems. Activation of the ATG1 complex in the initiation stage results in phosphorylation of ATG6 (Beclin1) of the PtdIns3K complex. This leads to its recruitment to the omegasome. The recruitment process is also supported by Vmp1, which accumulates in subdomains of the ER. The signal lipid PtdIns3P is generated by the PtdIns3K complex and is required for binding of other autophagy proteins to the membrane such as ATG18 and ATG2. The balance between production and degradation of PtdIns3P regulates omegasome formation and elongation of the isolation membrane and the transmembrane protein ATG9 is thought to deliver the required membrane lipids. For efficient membrane elongation two ubiquitin-like conjugation reactions are crucial. The ATG12~5/16 complex regulates the conjugation of ATG8 (LC3 in mammals) to phosphatidylethanolamine (PE) at the isolation membrane. ATG8b (blue) joins the growing autophagosome before ATG8a (red). Furthermore, ATG8b is mainly localised at the outside and ATG8a at the inside of the autophagosome. In the final steps ATG8a and b at the outer membrane are cleaved by ATG4 (not shown), the autophagosome fuses with a lysosome and the contents of the autophagosome are degraded in the autolysosome. The DDB_G numbers of the likely *D. discoideum* orthologues of the following mammalian proteins are provided: FIP200 (DDB_G0285767), ATG14 (DDB_G0283825), UVRAG (DDB_G0288175) and Bif-1 (DDB_G0274805). Components of the autophagic machinery are not drawn to scale. PI = phosphatidylinositol; PI3P = phosphatidylinositol-3-phosphate; PE = phosphatidyl-ethanolamine. Modified from (Mesquita et al., 2017).



tors (Farre and Subramani, 2016). They tether the cargo to the autophagosomal membrane by simultaneously binding the cargo and ATG8 family proteins (LC3 in mammals) present on the surface of the growing autophagosome (Zaffagnini and Martens, 2016). The interaction is mediated by different receptors, such as e.g. p62/SQSTM1, which contain ubiquitin-binding domains for the cargo and an LC3-interacting region (LIR), also called ATG8-interacting motif (AIM) (Boya et al., 2013, Gatica et al., 2018, Khaminets et al., 2016). Based on the cargo to be destroyed, selective autophagy is subdivided into ribophagy, mitophagy, aggrephagy, xenophagy, lipophagy, reticulophagy, nucleophagy, glycophagy, pexophagy, and proteophagy (Marshall and Vierstra, 2018).

***Dictyostelium discoideum* an excellent model system for autophagy**

Autophagy is intensively studied in vertebrates, e.g. *Mus*

musculus, insects, e.g. *Drosophila melanogaster*, worms, e.g. *Caenorhabditis elegans*, fungi, e.g. *S. cerevisiae* and plants, e.g. *Arabidopsis thaliana*, to name just a few (Galluzzi et al., 2017, Lv et al., 2014, Mesquita et al., 2017). Another of the well-established model organisms is the soil-living amoeba *D. discoideum* and in recent years, a large number of methods to monitor and quantify autophagy in this organism have been developed (Calvo-Garrido et al., 2010, Domínguez-Martín et al., 2017, Klionsky et al., 2016). *D. discoideum* is a member of the phylum Amoebozoa and was first isolated and described by Kenneth Raper (Raper, 1935). *Dictyostelium* cells grow as unicellular amoebae that divide by binary cell fission and feed on bacteria by phagocytosis (Kessin, 1981). Upon depletion of the food source, up to 100,000 solitary amoebae aggregate by chemotaxis towards cAMP. The cell aggregate (or pseudoplasmodium) differentiates via distinct morphological states into a mature fruiting body, composed of a mass of spore cells supported by a thin, long stalk made of vacuolised dead cells

(Eichinger, 2003).

Since development takes place in the absence of external nutrients, *D. discoideum* cells must mobilise a large fraction of the required energy for biosynthetic needs and morphogenesis by autophagy and glycogenolysis (Mesquita *et al.*, 2017). In addition, autophagy is also required for signalling pathways relevant for the developmental process (Calvo-Garrido *et al.*, 2010). For example, the formation of fruiting bodies, which mainly consist of viable spores and vacuolised, cellulose-walled, dead stalk cells, requires autophagy (Mesquita *et al.*, 2017). For the differentiation from vegetative cells into stalk cells by autophagic cell death (ACD) at least two distinct stimuli are necessary. The first stimulus is starvation together with cAMP to induce autophagy and the second required stimulus is the main stalk cell differentiation-inducing factor DIF-1, a small dichlorinated molecule (Giusti *et al.*, 2009, Mesquita *et al.*, 2017). A very recent report showed that in the absence of the core autophagy proteins ATG5, ATG7 or ATG9 vacuolization of stalk cells still takes place, suggesting that ACD is not dependent on canonical autophagy but may still depend on certain non-canonical autophagy (Yamada and Schaap, 2019). Moreover, the unconventional secretion of AcbA, the precursor of the signalling peptide SDF-2 (spore differentiation factor 2), is indispensable for spore formation and depends on autophagy (Duran *et al.*, 2010). As a consequence, autophagy malfunction generally results in reduced cell survival upon nitrogen starvation and in developmental abnormalities in the affected strains, which also supports the identification of autophagy-deficient mutants in the laboratory (Fischer *et al.*, 2019, King *et al.*, 2013, Mesquita *et*

al., 2017, Otto *et al.*, 2003). The developmental phenotypes range from a complete lack of aggregation to the formation of extremely small and crippled fruiting bodies with drastically reduced spore viability (Calvo-Garrido *et al.*, 2010, Fischer *et al.*, 2019, Mesquita *et al.*, 2015, Messling *et al.*, 2017, Otto *et al.*, 2003, Otto *et al.*, 2004, Tung *et al.*, 2010, Xiong *et al.*, 2015). Differences in the importance of the respective autophagy proteins for the functioning of autophagy and additional non-autophagic functions for some of these proteins are likely responsible for the diverse observed phenotypes (Fischer *et al.*, 2019). Indeed, in recent years more and more non-autophagic functions of core autophagy proteins have been reported (Malhotra *et al.*, 2015, Mauthe *et al.*, 2016, Nam *et al.*, 2017, Xiong *et al.*, 2015). Recent excellent reviews have covered the study of autophagy in *D. discoideum* in the contexts of mechanical stress, human disease, infection with pathogens and cell death pathways (Calvo-Garrido *et al.*, 2010, King *et al.*, 2011, Mesquita *et al.*, 2017). Furthermore, currently established methods have been exquisitely summarized (Domínguez-Martín *et al.*, 2017, Mesquita *et al.*, 2013). In this review, we focus on the general features and mechanisms of autophagy, autophagy-dependent and -independent roles of ATG proteins, and the crosstalk between autophagy and the ubiquitin-proteasome system (UPS).

General features and mechanisms of autophagy

The autophagic process is tightly controlled by the cell's nutritional and energy level and can be subdivided into initiation, maturation, and lysosomal degradation phases. The responsible

TABLE 1

MAIN AUTOPHAGY-RELATED PROTEINS IN A SELECTION OF MODEL ORGANISMS

<i>Homo sapiens</i>	<i>Dictyostelium discoideum</i>	<i>Saccharomyces cerevisiae</i>	<i>Drosophila melanogaster</i>	<i>Caenorhabditis elegans</i>
ATG1 protein complex subunits				
ULK1, ULK2	ATG1	ATG1	ATG1	UNC-51
ATG13	ATG13	ATG13	ATG13	ATG13
ATG101	ATG101	–	ATG101	EPG-9
FIP200	DDB_G0285767	ATG11, ATG17	ATG17	EPG-7
PtdIns3K protein complex subunits				
Beclin-1	ATG6	VPS30	ATG6	BEC-1
PIK3C3	VPS34	VPS34	PI3K59F	VPS-34
PIK3R4	VPS15	VPS15	VPS15	VPS-15
UVRAG	DDB_G0288175	VPS38	UVRAG	T23G11.7, Y34B4A.2
Bif-1	DDB_G0274805 (<i>ibrA</i>)	–	Endophilin B	F35A5.8
ATG14	DDB_G0283825	ATG14	ATG14	EPG-8
Ubiquitin-like conjugation systems				
ATG3	ATG3	ATG3	ATG3	ATG-3
ATG4A, 4B, 4C, 4D	ATG4	ATG4	ATG4A, 4B	ATG-4.1, 4.2
ATG5	ATG5	ATG5	ATG5	ATG-5
ATG7	ATG7	ATG7	ATG7	ATG-7
GABARAP, GABARAPL1, L2, L3	ATG8b	ATG8	ATG8a	LGG-1
MAP1LC3A, B, C	ATG8a	–	<i>ATG8b</i>	LGG-2
ATG10	ATG10	ATG10	ATG10	ATG-10
ATG12	ATG12	ATG12	ATG12	LGG-3
ATG16L1, L2	ATG16	ATG16	ATG16	ATG-16.1, 16.2
Membrane delivery system				
ATG2A, 2B	DDB_G0277419	ATG2	ATG2	ATG-2
ATG9A, 9B	ATG9	ATG9	ATG9	ATG-9
WIPI1, 2, 3 (Wdr45), WIPI4 (Wdr45)	ATG18, Wdr45l	ATG18	ATG18a, 18b	ATG-18
Vmp1	Vmp1	–	Tango5	EPG-3

Standalone letters and numbers refer to the protein mentioned before; –, currently no obvious corresponding orthologue; italic, uncertain grouping.

core machinery for autophagosome formation comprises more than 20 proteins, that are evolutionary highly conserved and engaged in different molecular complexes (Fig. 1; Table 1) (Birgisdottir *et al.*, 2013, Calvo-Garrido *et al.*, 2010, Feng *et al.*, 2014, Ktistakis and Tooze, 2016). In mammalian and *D. discoideum* cells, several autophagosomes can be generated simultaneously at multiple cytoplasmic sites, whereas this occurs only at a single spot near the vacuole in *S. cerevisiae* and it has been suggested that yeast is “the odd one out” (King, 2012, Mesquita *et al.*, 2017). The exact origin of the autophagosomal membrane remains an enigma after more than 20 years of investigation. Recent observations indicate that the endoplasmic reticulum (ER) in interplay with the Golgi, endosomes, the plasma membrane and mitochondria serve as membrane source in mammalian cells (Wei *et al.*, 2018). The initial structure is a *de novo* generated subdomain of the ER, the so-called omegasome (PAS, phagophore assembly site or preautophagosomal structure in *S. cerevisiae*), which then becomes the isolation membrane (IM, phagophore in *S. cerevisiae*) (Axe *et al.*, 2008).

The nutrient sensor target of rapamycin (TOR), a member of the phosphatidylinositol kinase-related family of serine/threonine protein kinases, receives and integrates extra- and intracellular signals, which mirror the energy and nutrient status of the cell (Calvo-Garrido *et al.*, 2010, Jung *et al.*, 2010, Noda, 2017). In its active form the mammalian TOR complex 1 (TORC1), composed of TOR, Raptor, Lst8 and Deptor, is the main inhibitor of autophagy. TORC1, the ATG1 and PtdIns3K complexes are well conserved in *D. discoideum* and evolutionarily close to the corresponding complexes in higher eukaryotes. Therefore, we think it is justified to infer that the corresponding activities, that were uncovered in different model organisms and are described in the following part, are also conserved in *D. discoideum*. In general, the activity of TORC1 is inhibited under starvation conditions. This results in rapid dephosphorylation of ATG13 and thereupon activation of the ATG1 complex (ULK1 in mammals), which consists of the ATG1 kinase, ATG13, ATG101 and the scaffold protein FIP200 (also known as RB1CC1 or ATG17; Fig. 1; Table 1). Interaction studies have demonstrated that ATG13 binds to both ATG1 and ATG101 and the latter in turn stabilises ATG13 (Mercer *et al.*, 2009, Mesquita *et al.*, 2015). The likely FIP200 homolog in *D. discoideum* is DDB_G0285767 (Li *et al.*, 2014, Mesquita *et al.*, 2017). The active ATG1 complex stimulates autophagy through phosphorylation of Beclin-1 (ATG6) in association with ATG14 of the class III phosphoinositide 3-kinase (PtdIns3K) complex (Park *et al.*, 2018). This leads to the generation of the phospholipid PtdIns3P by Vps34/PtdIns3K at the omegasome. The balance between production and degradation of PtdIns3P is pivotal for recruitment of further core autophagy proteins to the omegasome, such as the PtdIns3P binding proteins ATG18 (WIP1 in mammals) and ATG2 (Lindmo and Stenmark, 2006, Obara *et al.*, 2008). Recently it was shown, that ATG2 is a multifunctional protein that tethers membranes and acts as a lipid-transfer protein. As ATG2 localises to the contact site between the enlarging IM and the ER it was suggested that ATG2 transfers phospholipids from the ER exit sites (ERES) to the IM during the process of autophagosome formation (Osawa and Noda, 2019, Valverde *et al.*, 2019). Furthermore, the highly conserved transmembrane proteins ATG9 and Vmp1 are needed for omegasome formation, PtdIns3P signalling and membrane delivery in *D. discoideum* and mammalian cells (Calvo-Garrido *et*

al., 2010, Calvo-Garrido *et al.*, 2014, Mesquita *et al.*, 2017, Tábara *et al.*, 2018). ATG9 is the only known integral membrane protein of the core autophagy machinery and resides in small vesicles that are involved in the delivery of membrane lipids to the growing autophagosome (Xie and Klionsky, 2007). Its knock-out resulted in a pleiotropic phenotype in *D. discoideum* (Tung *et al.*, 2010). In Vmp1-deficient *D. discoideum* cells PtdIns3P production and the subsequent recruitment of the autophagy machinery to the ER is intact, however, the autophagic flux is blocked. The ER-resident protein Vmp1 seems to generate an ideal ER microenvironment required for the correct structure of the omegasome, which allows the IM to elongate and to become a functional autophagosome (Calvo-Garrido *et al.*, 2010, Mesquita *et al.*, 2017, Tábara *et al.*, 2018). Further regulatory components of the PtdIns3K complex are the myristoylated protein kinase Vps15 (PIK3R4 in mammals), UVRAG and Bif-1 (Table 1).

For the expansion of the IM eight highly conserved core autophagy proteins are indispensable (Fig. 1; Table 1) (Geng and Klionsky, 2008). These proteins are involved in two ubiquitin-like conjugation reactions and similar to the ubiquitin system, the two ubiquitin-like proteins ATG12 and ATG8 (LC3 in mammals) are finally attached to their substrate via this enzymatic pathway. In the first ubiquitin-like reaction, ATG12 is activated by the E1-like enzyme ATG7 and then conjugated to the E2-like enzyme ATG10 (Geng and Klionsky, 2008). Subsequently, ATG12 is covalently attached to its target protein ATG5, and two ATG12~5 conjugates in turn associate non-covalently with an ATG16 dimer and form a hetero-tetrameric complex (Mizushima *et al.*, 1999). The ATG12~5 conjugation reaction seems to be irreversible, since so far no enzyme for the cleavage of the isopeptide bond between ATG12 and ATG5 has been identified (Geng and Klionsky, 2008). In the final step of the second ubiquitin-like reaction, ATG8 (LC3) is reversibly attached to the lipid phosphatidylethanolamine (PE) on the expanding autophagosomal membrane via the E3-like activity of the ATG12~5/16 complex. It is believed that the complex brings the ATG8-carrying E2-like enzyme ATG3 in proximity to PE and determines the exact site of ATG8-PE conjugation on the autophagosomal membrane (Fujita *et al.*, 2008, Sakoh-Nakatogawa *et al.*, 2013, Walczak and Martens, 2013). ATG8-PE is present on both the inner and outer membranes of the IM. In mammals there are seven paralogues of the single yeast ATG8. Three are grouped into the MAP1-LC3 (microtubule-associated protein 1 – light chain 3) and four into the GABARAP/GATE16 (Gamma-aminobutyric acid receptor-associated protein/Golgi-associated ATPase enhancer of 16 kDa) subfamily (Shpilka *et al.*, 2011). *D. discoideum* harbours two paralogues, ATG8a and ATG8b, which have distinct functions in canonical autophagy. It was shown that they associate to the autophagosome in succession and that ATG8b mainly localises at the outer membrane of the autophagosome while ATG8a is mainly present at the inner membrane (Matthias *et al.*, 2016, Messling *et al.*, 2017). This differential localisation supports a more prominent role for ATG8b as an adapter for the autophagy machinery and in autophagosome lysosome fusion while ATG8a appears to function mainly as a binding partner for autophagy receptors. On the basis of function and localisation it was inferred that ATG8b is likely the *D. discoideum* orthologue of the GABARAP and ATG8a of the LC3 subfamily in mammals (Table 1) (Messling *et al.*, 2017). The autophagosomal membrane further expands through the incorporation of membrane lipids, engulfs entire organelles or

parts of the cytoplasm and finally closes into a double-membrane structure, the autophagosome (Lamb *et al.*, 2013, Stanley *et al.*, 2014). After completion of autophagosome biogenesis and before fusion with the lysosome, ATG8 is cleaved from the outer membrane by ATG4, while ATG8 on the inner membrane is degraded inside the lysosome (Geng and Klionsky, 2008, Kirisako *et al.*, 2000). Autophagosomes eventually mature into autolysosomes upon fusion of the outer autophagosomal membrane with the lysosomal membrane. Finally, the inner autophagosomal membrane and the cargo are degraded by lysosomal hydrolases (Fig. 1) (Wirawan *et al.*, 2012). The molecular mechanism of autophagosome-lysosome fusion has so far not been investigated in *D. discoideum*. Results from different experimental systems showed, that it consists of two phases: the autophagosome migration phase and the fusion phase. In the migration phase, transport of autophagosomes to the location of the lysosomes in the perinuclear region occurs along microtubules in a dynein-dependent manner (Jahreiss *et al.*, 2008). In the autophagosome-lysosome fusion step, three sets of protein families are involved, Rab GTPases, membrane tethering complexes and soluble Nethylmaleimide-sensitive-factor attachment protein receptors (SNAREs) (Nakamura and Yoshimori, 2017).

Autophagy-independent roles of ATG proteins

There is growing evidence in the literature for unconventional roles of ATG proteins, besides their function in canonical autophagy. RNA_{Seq} analysis of different mammalian cells in combination with a siRNA screen revealed that up to 36% of the autophagy-related genes encode proteins with additional unconventional functions (Mauthe *et al.*, 2016). For example in mouse embryonic fibroblasts, LC3-coated vesicles that differ from autophagosomes are necessary for the disposal of the ERAD effector protein EDEM1 in the process of ER-associated degradation (ERAD) (Cali *et al.*, 2008). Independent from its function in autophagy mammalian ATG7 has been implicated in nutrient deprivation induced cell cycle arrest via direct interaction and induction of p53 (Lee *et al.*, 2012). In addition, autophagy-independent functions have been reported in different organisms for ATG1, ATG2, ATG3, ATG4, ATG5, ATG6, ATG12, ATG16 and PtdIns3K (Bestebroer *et al.*, 2013, Schaaf *et al.*, 2016, Subramani and Malhotra, 2013). This list is likely not exhaustive and also in *Dictyostelium*, many of the core autophagy proteins appear to fulfil autophagy-independent functions in addition to their role in canonical autophagy (Xiong *et al.*, 2019, Xiong *et al.*, 2015).

LC3-associated phagocytosis

LC3-associated phagocytosis (LAP) is one such novel function for autophagy proteins and is a contributor to immune regulation and inflammatory responses across various cell and tissue types (Heckmann and Green, 2019). In contrast to canonical autophagy, LAP is not dependent on the AMPK–mTORC1–ULK1 (ATG1) activation axis or nutrient status of the cells (Heckmann *et al.*, 2017, Martinez *et al.*, 2015). A plurality of ligands, including dying cells, immune complexes and pathogens, has been shown to facilitate the conjugation of LC3 to PE of the phagosome in this process. Several receptors that participate in cargo recognition have already been identified including toll-like receptors, immunoglobulin receptors or TIM4 (Heckmann and Green, 2019). Following an activating stimulus, LAP can be delineated into three phases, followed by lysosomal fusion. The first phase is the generation of the single


membrane phagosome which serves as scaffold for the assembly of downstream regulatory complexes (Heckmann *et al.*, 2017). Secondly, the PtdIns3K complex composed of Vps34 (PtdIns3K), Vps15, Beclin-1 (ATG6 in e.g. yeast and *D. discoideum*), UVRAG and Rubicon generates membrane-localised PtdIns3P (Martinez *et al.*, 2015). Thirdly, the conjugation systems are recruited, which catalyse the conjugation of LC3 to PE of the phagosome (Fig. 2). The presence of LC3 on the phagosome, now termed LAPosome, mediates the fusion with the lysosome and the cargo is degraded (Heckmann and Green, 2019). LAP exists at the crossroads of phagocytosis and autophagy and is for example required for apoptotic corpse clearance during programmed cell death in multicellular organisms (Huang *et al.*, 2013, Martinez *et al.*, 2011). There is general agreement that LAP requires the entire ATG12~5/16 complex for LC3 recruitment (Lai and Devenish, 2012). Loss of any of the components of this complex led to a reduction or even abolishment of LAP function in mouse embryonic fibroblasts (MEFs) and mouse bone marrow-derived macrophages (Huang *et al.*, 2009, Kageyama *et al.*, 2011, Lai and Devenish, 2012). Investigation of the phagocytic and macropinocytic activity of the *D. discoideum* ATG9⁻, ATG16⁻, ATG9^{7/16}⁻, ATG12⁻ and ATG12^{7/16}⁻ strains showed that the encoded proteins have functions in phagocytosis of bacteria and yeast and also in the uptake of nutrients via macropinocytosis (Fischer *et al.*, 2019, Tung *et al.*, 2010, Xiong *et al.*, 2015). However, this is only circumstantial evidence and thus far LAP has not been directly demonstrated in *D. discoideum*. There could also be an indirect connection as for example Vmp1 deficiency in *Dictyostelium* led not only to a simple block in autophagosome formation but rather to a disturbance in autophagy-dependent PtdIns3P signalling at the ER, which also indirectly influenced macropinocytosis (Calvo-Garrido *et al.*, 2014). Electron microscopy in combination with immunogold labelling could resolve, whether there is LAP also in *Dictyostelium*.

Mammals express two ATG16 isoforms of which ATG16L1 apparently plays a critical role in the defence against pathogens. By recruiting ATG16L1 to the bacterial entry site, bacterial sensing by NOD proteins is linked to the induction of autophagy (Travassos *et al.*, 2010). Furthermore, ATG16L1 is required for LAP as murine bone marrow-derived macrophages deficient in ATG16L1 failed to undergo both, canonical autophagy or LAP (Martinez *et al.*, 2015). More recently, it was reported that the C-terminal domain of ATG16L1, which is composed of seven WD40 repeats, is essential for LAP during non-canonical autophagy, but dispensable for canonical autophagy (Fletcher *et al.*, 2018). This finding opens the possibility for the detailed analysis of LC3 lipidation during LAP.

Further autophagy-independent functions of ATG5, ATG12 and ATG16

In silico analyses showed that *D. discoideum*, yeast and human ATG5 harbour one helix-rich domain and two ubiquitin-like domains and ATG12 one ubiquitin-like domain. Although ubiquitin, ATG5, and ATG12 do not share sequence similarity, the 3D structures of the ubiquitin-like domains are highly similar, especially with respect to the α -helices and β -strands (Fischer *et al.*, 2019, Geng and Klionsky, 2008, Tsukada and Ohsumi, 1993). *D. discoideum* ATG16 is composed of three distinct regions, as is the case for the ATG16 orthologs in higher eukaryotes: the N-terminal domain which is responsible for binding to ATG5, followed by a coiled-coil domain (CCD) which mediates homo-dimerisation and seven WD40

Fig. 2. Mechanism of LC3-associated phagocytosis (LAP). LAP is characterised by employing parts of the canonical autophagy machinery for the conjugation of LC3 family proteins to the phagosomal membrane. It is initiated by engagement of specific receptors that recognise a variety of cargos including pathogens, dying cells or soluble ligands. The cargo is internalized by receptor-mediated phagocytosis and engulfed within a single-membrane phagosome. This compartment is rapidly decorated with phosphatidylinositol-3-phosphate (PtdIns3P) generated by the PtdIns3K complex consisting of Vps15, Vps34, Beclin-1 (ATG6), UVRAG and Rubicon. Autophagy proteins including the ATG12~5/16 complex are subsequently recruited to the surface of phagosomal membrane, which in turn covalently link LC3 (ATG8) to phosphatidylethanolamine (PE) on the surface of the phagosome. The presence of LC3 on the phagosome, now termed LAPosome, mediates fusion with the lysosome and the cargo is degraded. The DDB_G numbers of the likely *D. discoideum* orthologues of UVRAG and Rubicon are DDB_G0288175 and DDB_G0293570, respectively.



repeats in the C-terminal half of the protein, which are predicted to form a β -propeller structure (Xiong *et al.*, 2019). This structure serves as a hub for protein-protein interactions, which appear to be crucial for autophagy-dependent or -independent functions of ATG16 (Xiong *et al.*, 2019). The C-terminal domain is missing in yeast ATG16 and recently, the crystal structure of the yeast ATG12~5/16 complex has been solved, showing that ATG12 and ATG16 are located on opposite sides of ATG5 and that there is no molecular interaction between ATG16 and ATG12 (Suzuki *et al.*, 2017).

The activity of the conserved hetero-tetrameric ATG12~5/16 complex is indispensable for autophagosome formation. During canonical autophagy it localises to the outer membrane of the expanding IM and is released shortly before or after autophagosome completion (Mizushima *et al.*, 2001). The association of the ATG12~5 conjugate with ATG16 apparently unmask a membrane-binding site in ATG5 and the membrane tethering ability of ATG5 is also stimulated by ATG12 (Walczak and Martens, 2013). Within the ATG12~5/16 complex, ATG16 is required for correct localisation and the ATG12~5 conjugate possesses the E3 ligase activity that promotes the conjugation of ATG8 to PE of the autophagosomal membrane (Fujita *et al.*, 2008, Hanada *et al.*, 2007). Mice lacking ATG5, ATG12 or ATG16L1 survive the embryonic phase, but die one day after birth, corroborating the importance of an intact ATG12~5/16 complex for postnatal survival (Kuma *et al.*, 2004, Malhotra *et al.*, 2015, Saitoh *et al.*, 2008). With respect to functions in non-canonical autophagy the situation is complicated and it is often not clear whether these are mediated by ATG5, ATG12, ATG16, the ATG12~5 conjugate, ATG5/16 or the ATG12~5/16 complex. In MEFs the ATG12~5/16 complex contributed to pneumolysin toxin resistance and restriction of cell-to-cell spread of *Listeria monocytogenes* through a pathway independent of macroautophagy.

The authors showed that in MEFs deficient in ATG16L1, as well as ATG5 or ATG12, cholesterol accumulated in lysosomes. This resulted in less efficient lysosomal exocytosis, which is needed for efficient plasma membrane repair. Interestingly, MEFs harbouring the ATG16L1 T300A allele, which was previously found to increase the risk of Crohn's disease (CD), were also less efficient in plasma membrane repair (Serramito-Gómez *et al.*, 2016, Tan *et al.*, 2018). There is also evidence that ATG16L1 modulates the balance between NOD2-induced xenophagy versus cytokine production. This may explain the effects of this polymorphism on the inflammatory process in CD (Plantinga *et al.*, 2011). In addition, the autophagy machinery including components of the ubiquitin-like conjugation systems also has an important role in maintaining membrane integrity during mycobacterial infection (Cardenal-Muñoz *et al.*, 2017). In *D. discoideum*, membrane damages caused by *Mycobacterium marinum* activate an autophagic defence response reflected by an up-regulation of autophagy genes, stimulation of autophagosome formation and recruitment to the mycobacteria-containing vacuole (MCV). The autophagic flux is simultaneously repressed resulting in the accumulation of cytoplasmic material in the MCV, which supports bacterial survival within the niche (Cardenal-Muñoz *et al.*, 2017). On the other hand, the membrane generated by the autophagic machinery at the distal pole of the ejecting bacteria prevents plasma membrane leakage and cell death of *D. discoideum* cells (Gerstenmaier *et al.*, 2015). In *Drosophila*, ATG16 is crucial for the differentiation of intestinal stem cells into enteroendocrine (EE) cells. Expression of ATG16 lacking the WD40 repeat domain led to morphological changes in the intestine that resembled inflammatory bowel disease (Nagy *et al.*, 2017). Furthermore, the C-terminal WD40 domain is important in xenophagy (Xiong *et al.*, 2019). During infection of intestinal epithelial cells with *Salmonella typhimurium*, damage of the en-

dosomal membrane by bacteria residing in the endosome leads to the exposure of endosomal proteins, which are ubiquitinated. ATG16L1 is recruited to the *Salmonella*-containing endosomes by a direct interaction between the WD40 repeat domain and ubiquitin (Fujita *et al.*, 2013, Xiong *et al.*, 2019). ATG16 is also involved in Cullin-3-mediated ubiquitination and proteasomal degradation of the selective autophagy adaptor p62/SQSTM1 (Xiong *et al.*, 2019). In *Dictyostelium*, ATG16 carries out additional functions in proteasomal activity, axenic growth and macropinocytosis (Xiong *et al.*, 2018, Xiong *et al.*, 2019, Xiong *et al.*, 2015). Since the *Dictyostelium atg16* knockout mutant displayed strongly reduced proteasomal activity, ATG16 is crucial for optimal UPS function (Xiong *et al.*, 2015). Moreover, a strong defect of *Dictyostelium* ATG16⁻ cells in macropinocytosis, as evidenced by their reduced growth in liquid medium and reduced uptake of TRITC-labelled dextran, which could be caused by a disturbance of recycling endosomes, was observed (Fischer *et al.*, 2019, Xiong *et al.*, 2015).

Several autophagy-independent functions have been described for ATG12 in different organisms. Under certain nutrient-limiting conditions, ATG12 interacts with ATG3 in maintaining mitochondrial homeostasis and preventing cell death in MEFs, HeLa and HEK293 cells. In this process, ATG12 serves as a positive mediator of mitochondrial apoptosis and directly regulates the apoptotic pathway by binding and inactivating pro-survival Bcl-2 family members, including Bcl-2 and Mcl-1 (Radoshevich *et al.*, 2010, Rubinstein *et al.*, 2011). Furthermore, ATG12 is involved in endosomal trafficking and IFN γ -mediated host defence against murine norovirus (MNV) infection as shown in HeLa cells and bone marrow-derived macrophages (Hwang *et al.*, 2012, Murrow *et al.*, 2015). In viral protein translation, ATG12 is crucial for translation of hepatitis C RNA, virus replication and egress from cells (Dreux and Chisari, 2011). In *D. discoideum* we found massive transcriptional changes and complex phenotypes of varying severity for ATG12⁻, ATG16⁻ and ATG12⁻/16⁻ knock-out cells implying that ATG12 and ATG16 have, in addition to their role in canonical autophagy, autophagy-independent functions (Fischer *et al.*, 2019). The developmental phenotypes of the ATG12⁻, ATG16⁻ and ATG12⁻/16⁻ strains were similar to those previously reported for ATG5⁻, ATG7⁻ and ATG16⁻ mutants (Otto *et al.*, 2003, Xiong *et al.*, 2015). Loss of either of these proteins led to severe impairments in the tipped mound stage, in the slug stage, and in fruiting body formation. The ATG12⁻, ATG16⁻ and ATG12⁻/16⁻ strains displayed similar defects in autolysosome formation and cellular viability in response to amino acid starvation, implying that ATG12 and ATG16 act together with ATG5 as a functional unit in canonical autophagy. Interestingly, an incremental increase in the severity of the phenotype from ATG12⁻ to ATG16⁻ cells to the double knock-out strain was the case for spore viability and maximal cell titre in liquid culture. This suggests that ATG12 and ATG16 fulfil an independent function in these cellular processes or that the ATG12~5/16 complex without either ATG12 or ATG16 has still some residual activity. In contrast, for proteasomal activity, axenic growth, and macropinocytosis the phenotypes of ATG16⁻ cells were in comparison to ATG12⁻ cells slightly more severe. Therefore, ATG16 either fulfils an additional function in these cellular processes or there is still some ATG5/16 complex with residual activity formed in the absence of ATG12. Even though the exact link between autophagy and phagocytosis is currently not clear in *D. discoideum*, the results of the clearing assay for *K. aerogenes* suggest that ATG16 has an inhibitory and

ATG12 a stimulatory effect on the clearing of *K. aerogenes* by phagocytosis (Fischer *et al.*, 2019).

ATG5, as also is the case for ATG16L1, is important for the biology of the Paneth cells of the mouse ileal epithelium since ATG5-deficient Paneth cells have notable abnormalities in antimicrobial peptide secretion (Cadwell *et al.*, 2008). In osteoclasts, ATG5 is involved in the polarised secretion of lysosomal contents (DeSelm *et al.*, 2011). Moreover, ATG5 also participates in autophagy-independent defence mechanisms against the intracellular protozoan parasite *Toxoplasma gondii*. This parasite persists in the vacuole of non-activated macrophages by preventing the fusion of the modified parasitophorous vacuole, in which it replicates, with the lysosome. In particular, *T. gondii*-infected *atg5*-deficient macrophages are unable to clear the parasite (Bestebroer *et al.*, 2013). Apoptotic stimuli, like anticancer drugs stimulate the calpain-mediated ATG5 cleavage and the resulting truncated form of ATG5 associates with mitochondria and triggers cytochrome c release and subsequent caspase 3 activation. Therefore, ATG5 represents a molecular link between autophagy and apoptosis — a finding with potential importance for clinical anticancer therapies (Bestebroer *et al.*, 2013, Ye *et al.*, 2018).

The plurality of recent studies on autophagy-independent functions of ATG proteins emphasizes the importance of revisiting phenotypes and functions, that to date have been attributed to canonical autophagy based on the genetic analysis of a single individual ATG protein. Autophagy-related or -independent functions in different organisms include secretion, trafficking of phagocytosed material, replication and egress of viral particles, and regulation of inflammatory and immune signalling cascades. The multitude of autophagy-independent processes in diverse model organisms also clearly shows that the non-metazoan social amoeba *Dictyostelium* can only shed light on some of the multifaceted roles of ATG proteins in non-canonical autophagy. This exciting new facet of autophagy research deepens our understanding of autophagy-related functions and signalling pathways mediated by single ATG proteins as well as entire cellular processes using components of the autophagy machinery. Henceforth, it will be challenging to mechanistically separate autophagy from these related pathways.

Crosstalk between autophagy and the ubiquitin-proteasome system (UPS)

Until recently, the UPS and autophagy were considered to be two independent protein degradation machineries with no point of interaction, since both systems have different substrate preferences and separate molecular mechanisms (Korolchuk *et al.*, 2009, Korolchuk *et al.*, 2010). The proteasome is extremely efficient in degrading smaller short-lived, abnormal or damaged proteins, which require temporal control such as cell cycle-related proteins (Ciechanover and Kwon, 2015). In contrast larger long-lived proteins, protein aggregates or even whole organelles are delivered to and degraded in lysosomes via the autophagosomal route (Ciechanover and Kwon, 2015). However, the observation that more than 40 proteins are shared as either substrates or regulators of both autophagy and the UPS, among them core autophagy proteins like ATG5, ATG7, ATG8, and ATG16, has generated interest in analysing the crosstalk between these two pathways (Gao *et al.*, 2010, Komatsu *et al.*, 2005, Mizushima and Levine, 2010, Nam *et al.*, 2017, Xiong *et al.*, 2018). The most

crucial common denominator of both degradative pathways is ubiquitination as degradation signal on substrates (Kraft *et al.*, 2010). The mode of degradation is determined by the nature of ubiquitin chains, with mono-ubiquitinated substrates and K63-linked chains being preferably degraded by autophagy and K48-linked chains by the UPS (Kwon and Ciechanover, 2017). To connect ubiquitin and autophagy, selective autophagy requires ubiquitin-binding receptors such as p62/SQSTM1, NBR1 or HDAC6 (Schreiber and Peter, 2014). The strongest evidence for a functional interaction between the UPS and autophagy came from the finding that the autophagy pathway is activated in response to a decrease in UPS activity (Shen *et al.*, 2013). The up-regulation of autophagy genes is mainly mediated by the unfolded protein response (UPR) resulting in the activation of the transcription factor ATF4 and by mitochondria-originating ROS, which triggers autophagy through activated AMPK (B'Chir *et al.*, 2013, Zhao *et al.*, 2016). Whether autophagy inhibition triggers a compensatory increase of the UPS activity is still controversial. There have been contradictory reports describing compensatory up-regulation of the UPS, no change in UPS activity or even a decrease in the UPS activity (Korolchuk *et al.*, 2009, Liu *et al.*, 2016, Wang *et al.*, 2013). It must be noted that the majority of autophagosomal substrates are too large to be degraded through the cylindrical proteasome (Park and Cuervo, 2013). In recent years a number of studies reported that mice and *D. discoideum* strains lacking core autophagy genes, such as ATG5, ATG7, ATG8a, ATG12 and ATG16, showed an accumulation of ubiquitinated protein aggregates. In addition, the *D. discoideum* autophagy mutants displayed a significant decrease in their proteasomal activity in comparison to wild-type cells (Arhzaouy *et al.*, 2012, Fischer *et al.*, 2019, Komatsu *et al.*, 2005, Messling *et al.*, 2017, Mizushima and Levine, 2010, Xiong *et al.*, 2015). Interestingly, in mammals the half-life and cellular concentration of ATG8, ATG12 and ATG16 themselves appear to be regulated by the UPS and, *vice versa*, proteasomal subunits were found to be degraded by lysosomes (Cuervo *et al.*, 1995, Fujita *et al.*, 2009, Gao *et al.*, 2010, Haller *et al.*, 2014).

The close interconnection between the UPS and autophagy was further substantiated by the discovery of proteaphagy, a novel type of selective autophagy, in *S. cerevisiae*, HeLa cells and the mouse-ear cress *A. thaliana* (Cohen-Kaplan *et al.*, 2016, Marshall *et al.*, 2015, Marshall *et al.*, 2016, Waite *et al.*, 2016). In *A. thaliana*, the autophagic degradation of the 26S proteasome is mediated by the 19S proteasomal subunit RPN10 (PSMD4) via ubiquitin and ATG8 (Marshall *et al.*, 2015). In yeast the ubiquitin receptor Cue5 and the Hsp42 chaperone are responsible for degradation (Marshall *et al.*, 2016). In contrast, in mammals, stress-induced proteaphagy was dependent on p62/SQSTM1 and ubiquitin (Cohen-Kaplan *et al.*, 2016). It appears therefore that proteaphagy is universal in eukaryotes, however, the mechanism and the autophagy receptors vary from organism to organism. There is evidence in *D. discoideum* for a direct interaction of ATG16 with PSMD1 and PSMD2, two subunits of the 19S regulatory particle of the proteasome. It was also shown that PSMD1/2 get degraded in lysosomes. However, it is currently not clear how the interaction between ATG16 and PSMD1/2 is regulated and whether this interaction is responsible for proteaphagy in *D. discoideum* (Xiong *et al.*, 2018). Taken together, the data suggest that in autophagy-compromised *D. discoideum* strains less active or inactive proteasomes accumulate, which would cause the observed dramatic decrease in proteasomal

activity (Arhzaouy *et al.*, 2012, Fischer *et al.*, 2019, Messling *et al.*, 2017, Xiong *et al.*, 2015).

Conclusions

The “professional” phagocyte *D. discoideum* is an extremely suitable, non-mammalian model organism for the study of autophagy in the context of a whole organism due to its unique life style, with vegetative unicellular and multicellular stages. The haploid genome of *D. discoideum* is completely sequenced and annotated, the generation of single or multiple gene replacement mutants and of strains expressing tagged proteins is generally straightforward and a rich collection of biochemical, cell biological and microscopical methods is available to study mutant phenotypes. These advantages and the close similarity of its autophagic system to higher eukaryotes including man make *D. discoideum* a perfect system for the investigation of unanswered questions in canonical autophagy, for the analysis of autophagy-independent roles of ATG proteins and also for the discovery of new players in the fascinating autophagy world.

References

- ARHZAOUY, K., STRUCKSBERG, K.H., TUNG, S.M., TANGAVELOU, K., STUMPF, M., FAIX, J., SCHRODER, R., CLEMEN, C.S. and EICHINGER, L. (2012). Heteromeric p97/p97R155C complexes induce dominant negative changes in wild-type and autophagy 9-deficient *Dictyostelium* strains. *PLoS One* 7: e46879.
- AXE, E.L., WALKER, S.A., MANIFAVA, M., CHANDRA, P., RODERICK, H.L., HABERMANN, A., GRIFFITHS, G. and KTISTAKIS, N.T. (2008). Autophagosome formation from membrane compartments enriched in phosphatidylinositol 3-phosphate and dynamically connected to the endoplasmic reticulum. *J. Cell Biol.* 182: 685–701.
- B'CHIR, W., MAURIN, A.C., CARRARO, V., AVEROUS, J., JOUSSE, C., MURANISHI, Y., PARRY, L., STEPIEN, G., FAFOURNOUX, P. and BRUHAT, A. (2013). The eIF2alpha/ATF4 pathway is essential for stress-induced autophagy gene expression. *Nucleic Acids Res* 41: 7683–7699.
- BESTEBROER, J., V'KOVSKI, P., MAUTHE, M. and REGGIORI, F. (2013). Hidden behind autophagy: the unconventional roles of ATG proteins. *Traffic* 14: 1029–1041.
- BIRGISDOTTIR, A.B., LAMARK, T. and JOHANSEN, T. (2013). The LIR motif - crucial for selective autophagy. *J Cell Sci* 126: 3237–3247.
- BOYA, P., REGGIORI, F. and CODOGNO, P. (2013). Emerging regulation and functions of autophagy. *Nature Cell Biol.* 15: 713–720.
- CADWELL, K., LIU, J., BROWN, S., MIYOSHI, H., LOH, J., LENNERZ, J., KISHI, C., KC, W., CARRERO, J., HUNT, S. *et al.*, (2008). A key role for autophagy and the autophagy gene Atg16l1 in mouse and human intestinal Paneth cells. *Nature* 456: 259–263.
- CALI, T., GALLI, C., OLIVARI, S. and MOLINARI, M. (2008). Segregation and rapid turnover of EDEM1 by an autophagy-like mechanism modulates standard ERAD and folding activities. *Biochem Biophys Res Commun* 371: 405–410.
- CALVO-GARRIDO, J., CARILLA-LATORRE, S., KUBOHARA, Y., SANTOS-RODRIGO, N., MESQUITA, A., SOLDATI, T., GOLSTEIN, P. and ESCALANTE, R. (2010). Autophagy in *Dictyostelium*: Genes and pathways, cell death and infection. *Autophagy* 6: 686–701.
- CALVO-GARRIDO, J., KING, J.S., MUNOZ-BRACERAS, S. and ESCALANTE, R. (2014). Vmp1 regulates PtdIns3P signaling during autophagosome formation in *Dictyostelium discoideum*. *Traffic* 15: 1235–1246.
- CARDENAL-MUÑOZ, E., ARAFAH, S., LÓPEZ-JIMÉNEZ, A.T., KICKA, S., FALAISE, A., BACH, F., SCHAAD, O., KING, J.S., HAGEDORN, M. and SOLDATI, T. (2017). *Mycobacterium marinum* antagonistically induces an autophagic response while repressing the autophagic flux in a TORC1- and ESX-1-dependent manner. *PLoS Pathog* 13: 1–28.
- CIECHANOVER, A. and KWON, Y.T. (2015). Degradation of misfolded proteins in neurodegenerative diseases: therapeutic targets and strategies. *Exp Mol Med* 47: e147.

- COHEN-KAPLAN, V., LIVNEH, I., AVNI, N., COHEN-ROSENZWEIG, C. and CIECHANOVER, A. (2016). The ubiquitin-proteasome system and autophagy: Coordinated and independent activities. *Int J Biochem Cell Biol* 79: 403–418.
- CUERVO, A.M., PALMER, A., RIVETT, A.J. and KNECHT, E. (1995). Degradation of proteasomes by lysosomes in rat liver. *Eur J Biochem* 227: 792–800.
- DE DUVE, C. and WATTIAUX, R. (1966). Functions of Lysosomes. *Annu Rev Physiol* 28: 435–492.
- DESELM, C., MILLER, B., ZOU, W., BEATTY, W., VAN MEEL, E., TAKAHATA, Y., KLUMPERMAN, J., TOOZE, S., TEITELBAUM, S. and VIRGIN, H. (2011). Autophagy proteins regulate the secretory component of osteoclastic bone resorption. *Dev Cell* 21: 966–974.
- DOMÍNGUEZ-MARTÍN, E., CARDENAL-MUÑOZ, E., KING, J.S., SOLDATI, T., CORIA, R. and ESCALANTE, R. (2017). Methods to Monitor and Quantify Autophagy in the Social Amoeba *Dictyostelium discoideum*. *Cells* 6: 18.
- DREUX, M. and CHISARI, F.V. (2011). Impact of the autophagy machinery on hepatitis C virus infection. *Viruses* 3: 1342–1357.
- DURAN, J., ANJARD, C., STEFAN, C., LOOMIS, W. and MALHOTRA, V. (2010). Unconventional secretion of Acb1 is mediated by autophagosomes. *J Cell Biol* 188: 527–536.
- EICHINGER, L. (2003). Revamp a model-status and prospects of the *Dictyostelium* genome project. *Curr Genet* 44: 59–72.
- ESKELINEN, E.L. and SAFTIG, P. (2009). Autophagy: a lysosomal degradation pathway with a central role in health and disease. *Biochim Biophys Acta* 1793: 664–673.
- FARRE, J.C. and SUBRAMANI, S. (2016). Mechanistic insights into selective autophagy pathways: lessons from yeast. *Nat Rev Mol Cell Biol* 17: 537–552.
- FENG, Y., HE, D., YAO, Z. and KLIONSKY, D.J. (2014). The machinery of macroautophagy. *Cell Res* 24: 24–41.
- FISCHER, S., RIJAL, R., FROMMOLT, P., WAGLE, P., KONERTZ, R., FAIX, J., MESSLING, S. and EICHINGER, L. (2019). Functional Characterization of Ubiquitin-Like Core Autophagy Protein ATG12 in *Dictyostelium discoideum*. *Cells* 8: 1–27.
- FLETCHER, K., ULFERTS, R., JACQUIN, E., VEITH, T., GAMMOH, N., ARASTEH, J.M., MAYER, U., CARDING, S., WILEMAN, T., BEALE, R. *et al.*, (2018). The WD40 domain of ATG16L1 is required for its non-canonical role in lipidation of LC3 at single membranes. *EMBO J* 37: 17.
- FUJITA, N., ITOH, T., OMORI, H., FUKUDA, M., NODA, T. and YOSHIMORI, T. (2008). The Atg16L complex specifies the site of LC3 lipidation for membrane biogenesis in autophagy. *Mol Biol Cell* 19: 2092–2100.
- FUJITA, N., MORITA, E., ITOH, T., TANAKA, A., NAKAOKA, M., OSADA, Y., UMEMOTO, T., SAITOH, T., NAKATOGAWA, H., KOBAYASHI, S. *et al.*, (2013). Recruitment of the autophagic machinery to endosomes during infection is mediated by ubiquitin. *J Cell Biol* 203: 115–128.
- FUJITA, N., SAITOH, T., KAGEYAMA, S., AKIRA, S., NODA, T. and YOSHIMORI, T. (2009). Differential involvement of Atg16L1 in Crohn disease and canonical autophagy: analysis of the organization of the Atg16L1 complex in fibroblasts. *J Biol Chem* 284: 32602–32609.
- GALLUZZI, L., BAEHRECKE, E.H., BALLABIO, A., BOYA, P., BRAVO-SAN PEDRO, J.M., CECCONI, F., CHOI, A.M., CHU, C.T., CODOGNO, P., COLOMBO, M.I. *et al.*, (2017). Molecular definitions of autophagy and related processes. *EMBO J* 36: 1811–1836.
- GAO, Z., GAMMOH, N., WONG, P.-M., ERDJUMENT-BROMAGE, H., TEMPST, P. and JIANG, X. (2010). Processing of autophagic protein LC3 by the 20S proteasome. *Autophagy* 6: 126–137.
- GATICA, D., LAHIRI, V. and KLIONSKY, D.J. (2018). Cargo recognition and degradation by selective autophagy. *Nat Cell Biol* 20: 233–242.
- GENG, J. and KLIONSKY, D.J. (2008). The Atg8 and Atg12 ubiquitin-like conjugation systems in macroautophagy. Protein modifications: beyond the usual suspects. *EMBO Rep* 9: 859–864.
- GERSTENMAIER, L., PILLA, R., HERRMANN, L., HERRMANN, H., PRADO, M., VILLAFANO, G.J., KOLONKO, M., REIMER, R., SOLDATI, T., KING, J.S. *et al.*, (2015). The autophagic machinery ensures nonlytic transmission of mycobacteria. *Proc. Natl. Acad. Sci. USA* 112: 687–692.
- GIUSTI, C., TRESSE, E., LUCIANI, M.F. and GOLSTEIN, P. (2009). Autophagic cell death: analysis in *Dictyostelium*. *Biochim Biophys Acta* 1793: 1422–1431.
- HALLER, M., HOCK, A.K., GIAMPAZOLIAS, E., OBERST, A., GREEN, D.R., DEBNATH, J., RYAN, K.M., VOUSDEN, K.H. and TAIT, S.W. (2014). Ubiquitination and proteasomal degradation of ATG12 regulates its proapoptotic activity. *Autophagy* 10: 2269–2278.
- HANADA, T., NODA, N.N., SATOMI, Y., ICHIMURA, Y., FUJIOKA, Y., TAKAO, T., INAGAKI, F. and OHSUMI, Y. (2007). The Atg12-Atg5 conjugate has a novel E3-like activity for protein lipidation in autophagy. *J Biol Chem* 282: 37298–37302.
- HECKMANN, B.L., BOADA-ROMERO, E., CUNHA, L.D., MAGNE, J. and GREEN, D.R. (2017). LC3-Associated Phagocytosis and Inflammation. *J Mol Biol* 429: 3561–3576.
- HECKMANN, B.L. and GREEN, D.R. (2019). LC3-associated phagocytosis at a glance. *J Cell Sci* 132: 7.
- HUANG, J., CANADIEN, V., LAM, G.Y., STEINBERG, B.E., DINAUER, M.C., MAGALHAES, M.A., GLOGAUER, M., GRINSTEIN, S. and BRUMELL, J.H. (2009). Activation of antibacterial autophagy by NADPH oxidases. *Proc Natl Acad Sci U S A* 106: 6226–6231.
- HUANG, S., JIA, K., WANG, Y., ZHOU, Z. and LEVINE, B. (2013). Autophagy genes function in apoptotic cell corpse clearance during *C. elegans* embryonic development. *Autophagy* 9: 138–149.
- HWANG, S., MALONEY, N.S., BRUINSMA, M.W., GOEL, G., DUAN, E., ZHANG, L., SHRESTHA, B., DIAMOND, M.S., DANI, A., SOSNOVTSSEV, S.V. *et al.*, (2012). Nondegradative role of Atg5-Atg12/Atg16L1 autophagy protein complex in antiviral activity of interferon gamma. *Cell Host Microbe* 11: 397–409.
- JAHREISS, L., MENZIES, F.M. and RUBINSZTEIN, D.C. (2008). The itinerary of autophagosomes: from peripheral formation to kiss-and-run fusion with lysosomes. *Traffic* 9: 574–87.
- JUNG, C.H., RO, S.-H., CAO, J., OTTO, N.M. and KIM, D.-H. (2010). mTOR regulation of autophagy. *FEBS Lett* 584: 1287–1295.
- KAGEYAMA, S., OMORI, H., SAITOH, T., SONE, T., GUAN, J.L., AKIRA, S., IMAMOTO, F., NODA, T. and YOSHIMORI, T. (2011). The LC3 recruitment mechanism is separate from Atg9L1-dependent membrane formation in the autophagic response against Salmonella. *Mol Biol Cell* 22: 2290–2300.
- KESSIN, R.H. (1981). Conservatism in slime mold development. *Cell* 27: 241–243.
- KHAMINETS, A., BEHL, C. and DIKIC, I. (2016). Ubiquitin-Dependent and Independent Signals In Selective Autophagy. *Trends Cell Biol* 26: 6–16.
- KING, J.S. (2012). Autophagy across the eukaryotes - Is *S. cerevisiae* the odd one out? *Autophagy* 8: 1159–1162.
- KING, J.S., GUEHO, A., HAGEDORN, M., GOPALDASS, N., LEUBA, F., SOLDATI, T. and INSALL, R.H. (2013). WASH is required for lysosomal recycling and efficient autophagic and phagocytic digestion. *Mol Biol Cell* 24: 2714–2726.
- KING, J.S., VELTMAN, D.M. and INSALL, R.H. (2011). The induction of autophagy by mechanical stress. *Autophagy* 7: 1490–1499.
- KIRISAKO, T., ICHIMURA, Y., OKADA, H., KABEYA, Y., MIZUSHIMA, N., YOSHIMORI, T., OHSUMI, M., TAKAO, T., NODA, T. and OHSUMI, Y. (2000). The reversible modification regulates the membrane-binding state of Apg8/Aut7 essential for autophagy and the cytoplasm to vacuole targeting pathway. *J Cell Biol* 151: 263–276.
- KLIONSKY, D.J., ABDELMOHSEN, K., ABE, A., ABEDIN, M.J., ABELIOVICH, H., ACEVEDO, A.A., ADACHI, H., ADAMS, C.M., ADAMS, P.D., ADELI, K. *et al.*, (2016). Guidelines for the use and interpretation of assays for monitoring autophagy (3rd edition). *Autophagy* 12: 1–222.
- KOMATSU, M., WAGURI, S., UENO, T., IWATA, J., MURATA, S., TANIDA, I., EZAKI, J., MIZUSHIMA, N., OHSUMI, Y., UCHIYAMA, Y. *et al.*, (2005). Impairment of starvation-induced and constitutive autophagy in Atg7-deficient mice. *J Cell Biol* 169: 425–434.
- KOROLCHUK, V.I., MENZIES, F.M. and RUBINSZTEIN, D.C. (2009). A novel link between autophagy and the ubiquitin-proteasome system. *Autophagy* 5: 862–863.
- KOROLCHUK, V.I., MENZIES, F.M. and RUBINSZTEIN, D.C. (2010). Mechanisms of cross-talk between the ubiquitin-proteasome and autophagy-lysosome systems. *FEBS Lett* 584: 1393–1398.
- KRAFT, C., PETER, M. and HOFMANN, K. (2010). Selective autophagy: ubiquitin-mediated recognition and beyond. *Nat Cell Biol* 12: 836–841.
- KTISTAKIS, N.T. and TOOZE, S.A. (2016). Digesting the Expanding Mechanisms of Autophagy. *Trends Cell Biol* 26: 624–635.
- KUMA, A., HATANO, M., MATSUI, M., YAMAMOTO, A., NAKAYA, H., YOSHIMORI, T., OHSUMI, Y., TOKUHISA, T. and MIZUSHIMA, N. (2004). The role of autophagy during the early neonatal starvation period. *Nature* 432: 1032–1036.

- KWON, Y.T. and CIECHANOVER, A. (2017). The Ubiquitin Code in the Ubiquitin-Proteasome System and Autophagy. *Trends Biochem Sci* 42: 873–886.
- LAI, S. and DEVENISH, R.J. (2012). LC3-Associated Phagocytosis (LAP): Connections with Host Autophagy. *Cells* 1: 396–408.
- LAMB, C.A., YOSHIMORI, T. and TOOZE, S.A. (2013). The autophagosome: origins unknown, biogenesis complex. *Nat Rev Mol Cell Biol* 14: 759–774.
- LEE, I.H., KAWAI, Y., FERGUSSON, M.M., ROVIRA, I.I., BISHOP, A.J., MOTOYAMA, N., CAO, L. and FINKEL, T. (2012). Atg7 modulates p53 activity to regulate cell cycle and survival during metabolic stress. *Science* 336: 225–228.
- LEVINE, B. and KLIONSKY, D.J. (2017). Autophagy wins the 2016 Nobel Prize in Physiology or Medicine: Breakthroughs in baker's yeast fuel advances in biomedical research. *Proc Natl Acad Sci U S A* 114: 201–205.
- LI, Y., GAN, C.P., ZHANG, S., ZHOU, X.K., LI, X.F., WEI, Y.Q., YANG, J.L. and WU, M. (2014). FIP200 is Involved in Murine Pseudomonas Infection by Regulating HMGB1 Intracellular Translocation. *Cell Physiol Biochem* 33: 1733–1744.
- LINDMO, K. and STENMARK, H. (2006). Regulation of membrane traffic by phosphoinositide 3-kinases. *J Cell Sci* 119: 605–614.
- LIU, W.J., YE, L., HUANG, W.F., GUO, L.J., XU, Z.G., WU, H.L., YANG, C. and LIU, H.F. (2016). p62 links the autophagy pathway and the ubiquitin-proteasome system upon ubiquitinated protein degradation. *Cell Mol Biol Lett* 21: 29.
- LV, X., PU, X., QIN, G., ZHU, T. and LIN, H. (2014). The roles of autophagy in development and stress responses in *Arabidopsis Thaliana*. *Apoptosis* 19: 905–921.
- MALHOTRA, R., WARNE, J.P., SALAS, E., XU, A.W. and DEBNATH, J. (2015). Loss of Atg12, but not Atg5, in pro-opiomelanocortin neurons exacerbates diet-induced obesity. *Autophagy* 11: 145–154.
- MARSHALL, R.S., LI, F., GEMPERLINE, D.C., BOOK, A.J. and VIERSTRA, R.D. (2015). Autophagic Degradation of the 26S Proteasome Is Mediated by the Dual ATG8/Ubiquitin Receptor RPN10 in *Arabidopsis*. *Mol Cell* 58: 1053–1066.
- MARSHALL, R.S., M'CLOUGHLIN, F. and VIERSTRA, R.D. (2016). Autophagic Turnover of Inactive 26S Proteasomes in Yeast Is Directed by the Ubiquitin Receptor Cue5 and the Hsp42 Chaperone. *Cell Rep* 16: 1717–1732.
- MARSHALL, R.S. and VIERSTRA, R.D. (2018). Autophagy: The Master of Bulk and Selective Recycling. *Annu Rev Plant Biol* 69: 173–208.
- MARTINEZ, J., ALMENDINGER, J., OBERST, A., NESS, R., DILLON, C.P., FITZGERALD, P., HENGARTNER, M.O. and GREEN, D.R. (2011). Microtubule-associated protein 1 light chain 3 alpha (LC3)-associated phagocytosis is required for the efficient clearance of dead cells. *Proc Natl Acad Sci USA* 108: 17396–17401.
- MARTINEZ, J., MALIREDDI, R.K., LU, Q., CUNHA, L.D., PELLETIER, S., GINGRAS, S., ORCHARD, R., GUAN, J.L., TAN, H., PENG, J. *et al.*, (2015). Molecular characterization of LC3-associated phagocytosis reveals distinct roles for Rubicon, NOX2 and autophagy proteins. *Nat Cell Biol* 17: 893–906.
- MATTHIAS, J., MESSLING, S. and EICHINGER, L. (2016). The two *Dictyostelium* autophagy eight proteins, ATG8a and ATG8b, associate with the autophagosome in succession. *Eur. J. Cell Biol* 95: 15–25.
- MAUTHE, M., LANGEREIS, M., JUNG, J., ZHOU, X., JONES, A., OMTA, W., TOOZE, S.A., STORK, B., PALUDAN, S.R., AHOLA, T. *et al.*, (2016). An siRNA screen for ATG protein depletion reveals the extent of the unconventional functions of the autophagy proteome in virus replication. *J Cell Biol* 214: 619–635.
- MERCER, C.A., KALIAPPAN, A. and DENNIS, P.B. (2009). A novel, human Atg13 binding protein, Atg101, interacts with ULK1 and is essential for macroautophagy. *Autophagy* 5: 649–662.
- MESQUITA, A., CALVO-GARRIDO, J., CARILLA-LATORRE, S. and ESCALANTE, R. (2013). Monitoring autophagy in *Dictyostelium*. *Methods Mol Biol* 983: 461–470.
- MESQUITA, A., CARDENAL-MUNOZ, E., DOMINGUEZ, E., MUNOZ-BRACERAS, S., NUNEZ-CORCUERA, B., PHILLIPS, B.A., TABARA, L.C., XIONG, Q., CORIA, R., EICHINGER, L. *et al.*, (2017). Autophagy in *Dictyostelium*: Mechanisms, regulation and disease in a simple biomedical model. *Autophagy* 13: 24–40.
- MESQUITA, A., TABARA, L.C., MARTINEZ-COSTA, O., SANTOS-RODRIGO, N., VINCENT, O. and ESCALANTE, R. (2015). Dissecting the function of Atg1 complex in *Dictyostelium* autophagy reveals a connection with the pentose phosphate pathway enzyme transketolase. *Open Biol* 5: 150088.
- MESSLING, S., MATTHIAS, J., XIONG, Q., FISCHER, S. and EICHINGER, L. (2017). The two *Dictyostelium discoideum* autophagy 8 proteins have distinct autophagic functions. *Eur J Cell Biol* 96: 312–324.
- MIZUSHIMA, N. and LEVINE, B. (2010). Autophagy in mammalian development and differentiation. *Nat Cell Biol* 12: 823–830.
- MIZUSHIMA, N., LEVINE, B., CUERVO, A.M. and KLIONSKY, D.J. (2008). Autophagy fights disease through cellular self-digestion. *Nature* 451: 1069–1075.
- MIZUSHIMA, N., NODA, T. and OHSUMI, Y. (1999). Apg16p is required for the function of the Apg12p–Apg5p conjugate in the yeast autophagy pathway. *EMBO J* 18: 3888–3896.
- MIZUSHIMA, N., YAMAMOTO, A., HATANO, M., KOBAYASHI, Y., KABEYA, Y., SUZUKI, K., TOKUHISA, T., OHSUMI, Y. and YOSHIMORI, T. (2001). Dissection of autophagosome formation using Apg5-deficient mouse embryonic stem cells. *J Cell Biol* 152: 657–668.
- MURROW, L., MALHOTRA, R. and DEBNATH, J. (2015). ATG12-ATG3 interacts with Alix to promote basal autophagic flux and late endosome function. *Nat Cell Biol* 17: 300–310.
- NAGY, P., SZATMÁRI, Z., SÁNDOR, G.O., LIPPAI, M., HEGEDŰS, K. and JUHÁSZ, G. (2017). *Drosophila* Atg16 promotes enteroendocrine cell differentiation via regulation of intestinal Slit/Robo signaling. *Development* 144: 3990–4001.
- NAKAMURA, S. and YOSHIMORI, T. (2017). New insights into autophagosome-lysosome fusion. *J Cell Sci* 130: 1209–1216.
- NAM, T., HAN, J.H., DEVKOTA, S. and LEE, H.W. (2017). Emerging Paradigm of Crosstalk between Autophagy and the Ubiquitin-Proteasome System. *Mol Cells* 40: 897–905.
- NODA, T. (2017). Regulation of Autophagy through TORC1 and mTORC1. *Biomolecules* 7.
- OBARA, K., SEKITO, T., NIIMI, K. and OHSUMI, Y. (2008). The Atg18-Atg2 complex is recruited to autophagic membranes via phosphatidylinositol 3-phosphate and exerts an essential function. *J Biol Chem* 283: 23972–23980.
- OSAWA, T. and NODA, N.N. (2019). Atg2: A novel phospholipid transfer protein that mediates de novo autophagosome biogenesis. *Protein Sci* 28: 1005–1012.
- OTTO, G.P., WU, M.Y., KAZGAN, N., ANDERSON, O.R. and KESSIN, R.H. (2003). Macroautophagy is required for multicellular development of the social amoeba *Dictyostelium discoideum*. *J Biol Chem* 278: 17636–17645.
- OTTO, G.P., WU, M.Y., KAZGAN, N., ANDERSON, O.R. and KESSIN, R.H. (2004). *Dictyostelium* macroautophagy mutants vary in the severity of their developmental defects. *J Biol Chem* 279: 15621–15629.
- PARK, C. and CUERVO, A.M. (2013). Selective autophagy: talking with the UPS. *Cell Biochem Biophys* 67: 3–13.
- PARK, J.M., SEO, M., JUNG, C.H., GRUNWALD, D., STONE, M., OTTO, N.M., TOSO, E., AHN, Y., KYBA, M., GRIFFIN, T.J. *et al.*, (2018). ULK1 phosphorylates Ser30 of BECN1 in association with ATG14 to stimulate autophagy induction. *Autophagy* 14: 584–597.
- PLANTINGA, T.S., JOOSTEN, L.A. and NETEA, M.G. (2011). ATG16L1 polymorphisms are associated with NOD2-induced hyperinflammation. *Autophagy* 7: 1074–1075.
- RADOSHEVICH, L., MURROW, L., CHEN, N., FERNANDEZ, E., ROY, S., FUNG, C. and DEBNATH, J. (2010). ATG12 conjugation to ATG3 regulates mitochondrial homeostasis and cell death. *Cell* 142: 590–600.
- RAPER, K.B. (1935). *Dictyostelium discoideum*, a new species of slime mold from decaying forest leaves. *J Agr Res* 50: 135–147.
- RUBINSTEIN, A.D., EISENSTEIN, M., BER, Y., BIALIK, S. and KIMCHI, A. (2011). The autophagy protein Atg12 associates with antiapoptotic Bcl-2 family members to promote mitochondrial apoptosis. *Mol Cell* 44: 698–709.
- SAITOH, T., FUJITA, N., JANG, M.H., UEMATSU, S., YANG, B.G., SATOH, T., OMORI, H., NODA, T., YAMAMOTO, N., KOMATSU, M. *et al.*, (2008). Loss of the autophagy protein Atg16L1 enhances endotoxin-induced IL-1beta production. *Nature* 456: 264–268.
- SAKOH-NAKATOGAWA, M., MATOBA, K., ASAI, E., KIRISAKO, H., ISHII, J., NODA, N.N., INAGAKI, F., NAKATOGAWA, H. and OHSUMI, Y. (2013). Atg12-Atg5 conjugate enhances E2 activity of Atg3 by rearranging its catalytic site. *Nat Struct Mol Biol* 20: 433–439.
- SCHAAF, M.B., KEULERS, T.G., VOOIJS, M.A. and ROUSCHOP, K.M. (2016). LC3/GABARAP family proteins: autophagy-(un)related functions. *FASEB J* 30: 3961–3978.
- SCHNEIDER, J.L. and CUERVO, A.M. (2014). Autophagy and human disease: emerging themes. *Curr Opin Genet Dev* 26: 16–23.
- SCHREIBER, A. and PETER, M. (2014). Substrate recognition in selective autophagy and the ubiquitin-proteasome system. *Biochim Biophys Acta* 1843: 163–181.

- SERRAMITO-GÓMEZ, I., BOADA-ROMERO, E. and PIMENTEL-MUIÑOS, F.X. (2016). Unconventional autophagy mediated by the WD40 domain of ATG16L1 is derailed by the T300A Crohn disease risk polymorphism. *Autophagy* 12: 2254–2255.
- SHEN, Y.F., TANG, Y., ZHANG, X.J., HUANG, K.X. and LE, W.D. (2013). Adaptive changes in autophagy after UPS impairment in Parkinson's disease. *Acta Pharmacol Sin* 34: 667–673.
- SHPIILKA, T., WEIDBERG, H., PIETROKOVSKI, S. and ELAZAR, Z. (2011). Atg8: an autophagy-related ubiquitin-like protein family. *Genome Biol* 12: 226.
- STANLEY, R.E., RAGUSA, M.J. and HURLEY, J.H. (2014). The beginning of the end: how scaffolds nucleate autophagosome biogenesis. *Trends Cell Biol* 24: 73–81.
- SUBRAMANI, S. and MALHOTRA, V. (2013). Non-autophagic roles of autophagy-related proteins. *EMBO Rep* 14: 143–151.
- SUZUKI, H., OSAWA, T., FUJIOKA, Y. and NODA, N.N. (2017). Structural biology of the core autophagy machinery. *Curr Opin Struct Biol* 43: 10–17.
- TÁBARA, L.C., VICENTE, J.J., BIAZIK, J., ESKELINEN, E.L., VINCENT, O. and ESCALANTE, R. (2018). Vacuole membrane protein 1 marks endoplasmic reticulum subdomains enriched in phospholipid synthesizing enzymes and is required for phosphoinositide distribution. *Traffic* 19: 624–638.
- TAN, J., MELLOUK, N., OSBORNE, S., AMMENDOLIA, D., DYER, D., LI, R., BRUNEN, D., VAN RIJN, J., HUANG, J., CZUCZMAN, M. *et al.*, (2018). An ATG16L1-dependent pathway promotes plasma membrane repair and limits *Listeria monocytogenes* cell-to-cell spread. *Nat Microbiol* 3: 1472–1485.
- TRAVASSOS, L., CARNEIRO, L., RAMJEET, M., HUSSEY, S., KIM, Y., MAGALHÃES, J., YUAN, L., SOARES, F., CHEA, E., LE BOURHIS, L. *et al.*, (2010). Nod1 and Nod2 direct autophagy by recruiting ATG16L1 to the plasma membrane at the site of bacterial entry. *Nat Immunol* 11: 55–62.
- TSUKADA, M. and OHSUMI, Y. (1993). Isolation and characterization of autophagy-defective mutants of *Saccharomyces cerevisiae*. *FEBS Lett* 333: 169–174.
- TUNG, S.M., UNAL, C., LEY, A., PENA, C., TUNGGAL, B., NOEGEL, A.A., KRUT, O., STEINERT, M. and EICHINGER, L. (2010). Loss of *Dictyostelium* ATG9 results in a pleiotropic phenotype affecting growth, development, phagocytosis and clearance and replication of *Legionella pneumophila*. *Cell Microbiol* 12: 765–780.
- VALVERDE, D.P., YU, S., BOGGAVARAPU, V., KUMAR, N., LEES, J.A., WALZ, T., REINISCH, K.M. and MELIA, T.J. (2019). ATG2 transports lipids to promote autophagosome biogenesis. *J. Cell Biol.* 218 1787–1798.
- WAITE, K.A., DE-LA MOTA-PEYNADO, A., VONTZ, G. and ROELOFS, J. (2016). Starvation Induces Proteasome Autophagy with Different Pathways for Core and Regulatory Particles. *J Biol Chem* 291: 3239–3253.
- WALCZAK, M. and MARTENS, S. (2013). Dissecting the role of the Atg12-Atg5-Atg16 complex during autophagosome formation. *Autophagy* 9: 424–425.
- WANG, X.J., YU, J., WONG, S.H., CHENG, A.S., CHAN, F.K., NG, S.S., CHO, C.H., SUNG, J.J. and WU, W.K. (2013). A novel crosstalk between two major protein degradation systems: regulation of proteasomal activity by autophagy. *Autophagy* 9: 1500–1508.
- WEI, Y., LIU, M., LI, X., LIU, J. and LI, H. (2018). Origin of the Autophagosome Membrane in Mammals. *Biomed Res Int* 2018: 9.
- WIRAWAN, E., VANDEN BERGHE, T., LIPPENS, S., AGOSTINIS, P. and VANDE-NABEELE, P. (2012). Autophagy: for better or for worse. *Cell Res* 22: 43–61.
- XIE, Z. and KLIONSKY, D.J. (2007). Autophagosome formation: core machinery and adaptations. *Nat Cell Biol* 9: 1102–1109.
- XIONG, Q., FISCHER, S., KAROW, M., MÜLLER, R., MESSLING, S. and EICHINGER, L. (2018). ATG16 mediates the autophagic degradation of the 19S proteasomal subunits PSMD1 and PSMD2. *Eur J Cell Biol* 97: 523–532.
- XIONG, Q., LI, W., LI, P., YANG, M., WU, C. and EICHINGER, L. (2019). The Role of ATG16 in Autophagy and The Ubiquitin Proteasome System. *Cells* 8: 16.
- XIONG, Q., UNAL, C., MATTHIAS, J., STEINERT, M. and EICHINGER, L. (2015). The phenotypes of ATG9, ATG16 and ATG9/16 knock-out mutants imply autophagy-dependent and -independent functions. *Open Biol* 5: 150008.
- YAMADA, Y. and SCHAAP, P. (2019). Cyclic AMP induction of *Dictyostelium* prespore gene expression requires autophagy. *Dev Biol* 452: 114–126.
- YE, X., ZHOU, X. and ZHANG, H. (2018). Exploring the Role of Autophagy-Related Gene 5 (ATG5) Yields Important Insights Into Autophagy in Autoimmune/Autoinflammatory Diseases. *Front Immunol* 17: 2334.
- ZAFFAGNINI, G. and MARTENS, S. (2016). Mechanisms of Selective Autophagy. *J Mol Biol* 428: 1714–1724.
- ZHAO, B., QIANG, L., JOSEPH, J., KALYANARAMAN, B., VIOLLET, B. and HE, Y.Y. (2016). Mitochondrial dysfunction activates the AMPK signaling and autophagy to promote cell survival. *Genes Dis* 3: 82–87.

Further Related Reading, published previously in the *Int. J. Dev. Biol.*

Dictyostelium discoideum Sir2D modulates cell-type specific gene expression and is involved in autophagy

Rakhee Lohia, Punita Jain, Mukul Jain, Pradeep Kumar Burma, Anju Shrivastava and Shweta Saran

Int. J. Dev. Biol. (2017) 61: 95-104

<https://doi.org/10.1387/ijdb.160038ss>

The Dictyostelium prestalk inducer DIF-1 directs phosphorylation of a bZIP transcription factor

Yoko Yamada, Yuzuru Kubohara, Haruhisa Kikuchi, Yoshiteru Oshima, Hong-Yu Wang, Susan Ross and Jeffrey G. Williams

Int. J. Dev. Biol. (2013) 57: 375-381

<https://doi.org/10.1387/ijdb.130046jw>

Bimodal distribution of motility and cell fate in Dictyostelium discoideum

Pavana Goury-Sistla, Vidyanand Nanjundiah and Gopal Pande

Int. J. Dev. Biol. (2012) 56: 263-272

<https://doi.org/10.1387/ijdb.113384ps>

Dictyostelium discoideum: a model system for differentiation and patterning

R Escalante and J J Vicente

Int. J. Dev. Biol. (2000) 44: 819-835

<http://www.intjdevbiol.com/web/paper/11206323>

Cell-cell signaling and adhesion in phagocytosis and early development of Dictyostelium

E Bracco, B Pergolizzi, B Peracino, E Ponte, A Balbo, A Mai, A Ceccarelli and S Bozzaro

Int. J. Dev. Biol. (2000) 44: 733-742

<http://www.intjdevbiol.com/web/paper/11061438>

Development at the edge of multi-cellularity: Dictyostelium discoideum

R R Kay

Int. J. Dev. Biol. (2000) 44: 35-38

<http://www.intjdevbiol.com/web/paper/10761844>

Regulation of cell differentiation and pattern formation in Dictyostelium development

I Takeuchi, M Tasaka, K Okamoto and Y Maeda

Int. J. Dev. Biol. (1994) 38: 311-319

<http://www.intjdevbiol.com/web/paper/7981039>

